

## **12. Exploring Environmental Exposures of Breast Cancer Risk through Dietary**

**Zeliha Selamoglu**

Department of Medical Biology,  
Medicine Faculty, Nigde Omer Halisdemir University,  
Nigde, Turkey.

**Muhammad Yasir Naeem**

Department of Plant Production and Technologies,  
Faculty of Agricultural Sciences and Technologies,  
Nigde Omer Halisdemir University, Nigde, Turkey.

**Elifsena Canan Alp Arici**

Department of Obstetrics and Gynecology,  
Faculty of Medicine, Necmettin Erbakan,  
University of Meram, Konya, Turkey.

**Muhammad Raza Naqvi**

Medical Director (Hematologist/Oncologist) and  
Cancer Liason Physician Inter Mountain Health Care,  
Denver, USA.

**R. M. Madhumitha Sri.**

Department of Pharmaceutical Technology,  
Alagappa College of Technology, Anna University, Chennai.  
Associate Analyst, Zifo Technologies Pvt. Ltd.,  
Manapakkam, Chennai, Tamilnadu.

**Karthika P. R.**

Student, B.Tech. Aerospace Engineering,  
School of Mechanical Engineering,  
Lovely Professional University, Jalandhar (Punjab).

**S. Ravichandran**

Department of Chemistry,  
School of Mechanical Engineering, Lovely Professional university,  
Jalandhar, Punjab, India.



**Figure 12.1: Breast Cancer**

***Abstract:***

*Estrogen, a key sex hormone, is a known contributor to breast cancer risk. This review dissects the complex relationship between dietary and environmental estrogen exposures and breast cancer development. We explore diverse sources, encompassing plant-based estrogens (e.g., soy isoflavones) and environmental contaminants (e.g., bisphenol A). The review delves into the mechanisms by which these exposures might influence cancer risk, focusing on estrogen receptor interactions and downstream signaling pathways that promote breast cell proliferation and survival. While risk assessment tools exist, the review empowers individuals by highlighting modifiable lifestyle factors for potentially mitigating risk. We emphasize maintaining a healthy weight and engaging in regular physical activity. However, ongoing controversies and limitations in current research are acknowledged, particularly regarding the strength of associations between specific exposures and cancer risk. Finally, the review advocates for empowering individuals with evidence-based information. By fostering informed choices about diet, lifestyle, and potentially beneficial interventions, we can contribute to a future where women have a greater ability to manage their breast cancer risk profile.*

***Keywords:***

*Estrogen, breast cancer, dietary estrogens, environmental estrogens, risk assessment, prevention, lifestyle modifications, informed choices*

**12.1 Introduction:**

Breast cancer, the most prevalent non-skin cancer and leading cause of cancer death among women globally, casts a long shadow (Figure 12.1). The American Cancer Society estimates a staggering 288,200 new cases of invasive breast cancer to be diagnosed in women within the United States for 2024 alone. Understanding the risk factors associated with breast cancer development is paramount for early detection, preventative strategies, and the development of effective treatment options (Giaquinto et al., 2022). Estrogen, a critical female sex hormone, plays a well-defined role in breast cancer. It acts as a growth promoter for breast cells, and breast cancers that possess receptors for estrogen (estrogen receptor-

positive, ER+) rely heavily on its presence. This established knowledge has led to the successful development of hormone therapy treatments that either block estrogen receptors or reduce estrogen production (Miziak et al., 2023).

Beyond the naturally produced estrogen within the body, however, scientific inquiry is increasingly focused on the potential impact of exogenous estrogens on breast cancer risk. Exogenous estrogens are those we encounter from external sources, broadly categorized into dietary estrogens and environmental estrogens (Maharjan et al., 2021).

Dietary estrogens are naturally occurring compounds present in certain foods. Examples include phytoestrogens found in soy products and mycoestrogens produced by molds on grains. Environmental estrogens, also known as xenoestrogens, are man-made chemicals that mimic the effects of natural estrogen in the body. These chemicals permeate various everyday products, including plastics, pesticides, personal care products, and industrial processes (Paterni et al., 2017, Kinkad et al., 2021).

The potential link between these exogenous estrogens and breast cancer risk has sparked significant scientific investigation and public concern. This heightened awareness stems from two key factors:

**Ubiquitous Exposure:** Modern life exposes us to a constant influx of these external estrogens. Dietary choices, the materials we use, and even the environment we inhabit can contribute to our overall exposure. This ubiquity raises concerns about the potential cumulative impact on breast health over time.

**Estrogen's Role in Breast Cancer:** As mentioned earlier, estrogen's well-established role in promoting breast cell growth necessitates a closer look at any external sources that might mimic its effects. Understanding if and how these exogenous estrogens interact with the body's hormonal system becomes crucial for assessing their potential influence on breast cancer risk (Al-Shami et al., 2023, Gompel et al., 2020).

### **12.1.1 Aim of the Study:**

This review aims to delve into the current understanding of how dietary and environmental estrogenic exposures might influence breast cancer risk. We will explore the different types of these exposures, examine the mechanisms by which they might interact with the body, and analyze the evidence from research studies to understand the current state of knowledge and ongoing debates in this field.

However, before embarking on this deep dive, it's important to acknowledge the complexities involved in researching the link between exogenous estrogens and breast cancer. Here are some key challenges:

**Difficulties in Quantifying Exposure:** Accurately measuring an individual's overall exposure to dietary and environmental estrogens remains a challenge. Dietary intake varies, and environmental exposure depends on a multitude of factors like geographic location, occupation, and lifestyle choices.

Distinguishing Cause from Correlation: Observational studies often show associations between exposures and outcomes, but they cannot definitively prove causation. Other risk factors for breast cancer might confound the results, making it difficult to isolate the specific impact of exogenous estrogens. Diversity of Estrogenic Effects: Exogenous estrogens exhibit varying potencies and may even have anti-estrogenic properties in certain circumstances. Understanding the specific effects of different exposures adds another layer of complexity to the research.

Despite these challenges, ongoing research is shedding light on the potential influence of exogenous estrogens on breast cancer risk.

This review aims to present a comprehensive overview of this evolving field, highlighting the key findings, controversies, and areas for further investigation. By exploring this complex relationship, we can contribute to a more informed understanding of breast cancer risk and potentially identify new strategies for prevention.

## **12.2 Estrogenic Exposures: A Complex Landscape:**

Estrogenic exposures encompass a wide range of natural and man-made compounds that can mimic the effects of estrogen in the body. These exposures can be broadly categorized into two main groups: dietary estrogens and environmental estrogens (Gupta et al., 2020).

Understanding these exposures and their potential impact on breast cancer risk is crucial for a more comprehensive picture of breast health (Bilal et al., 2022).

### **12.2.1 Dietary Estrogens: A Natural Mix from Plants:**

Dietary estrogens are naturally occurring compounds found in various plant-based foods. While considerably weaker than endogenous estrogen, they can bind to estrogen receptors and exert weak estrogenic or anti-estrogenic effects depending on the specific compound and the overall hormonal milieu (Ionesc et al., 2021, Domínguez-López et al., 2020). Two main types of dietary estrogens are of particular interest:

**Phytoestrogens:** These are plant-derived compounds with structural similarities to estradiol, the most potent form of natural estrogen. Common dietary sources include soybeans and soy products (tofu, tempeh), legumes (lentils, chickpeas), flaxseeds, and certain grains (whole grains, oats) (Desmawati and Sulastrri, 2019).

The health effects of phytoestrogens are a subject of ongoing research. Some studies suggest potential benefits like reducing the risk of heart disease and menopausal symptoms. However, the impact on breast cancer risk is less clear. While some studies show no association, others suggest a possible protective effect at moderate intake levels (Rietjens et al., 2017).

**Mycoestrogens:** These are estrogenic compounds produced by molds that can grow on grains and other stored foods under warm and humid conditions. Zearalenone and deoxynivalenol (DON) are two commonly encountered mycoestrogens.

Exposure can occur through the consumption of contaminated grains or food products (Awuchi et al., 2022, Thapa et al., 2021). The potential health risks of mycoestrogens are of concern, with some evidence suggesting they might contribute to hormonal imbalances and even some cancers. However, the specific link to breast cancer risk remains inconclusive and requires further investigation (Thapa et al., 2021).

### **12.2.2 Environmental Estrogens: Xenoestrogens in Our Surroundings:**

Environmental estrogens, also known as xenoestrogens, are man-made chemicals that can mimic the effects of natural estrogen. They are ubiquitous in modern life and can be found in various products and environmental sources:

**Plastics:** Certain types of plastics, particularly those containing bisphenol A (BPA) or phthalates, can leach out xenoestrogens when heated or exposed to sunlight. These chemicals can then contaminate food and beverages stored in plastic containers (Zimmermann et al., 2021).

**Pesticides:** Some commonly used pesticides, such as DDT and atrazine, have been shown to exhibit estrogenic activity. Dietary exposure can occur through consumption of produce treated with these pesticides (Vandenberg et al., 2020).

**Personal Care Products:** Certain ingredients in cosmetics, lotions, and sunscreens have been identified as potential xenoestrogens. Absorption through the skin or inhalation can be potential routes of exposure (Vandenberg et al., 2020).

**Industrial Chemicals:** Specific chemicals used in industrial processes also exhibit estrogenic activity. Occupational exposure can be a concern for workers in related industries. The widespread presence of environmental estrogens raises concerns about their potential cumulative effects on human health, particularly breast cancer risk. However, research findings are mixed. Some studies suggest a possible link between exposure to certain xenoestrogens and an increased risk of breast cancer, while others haven't found a clear association (Rocha et al., 2021).

### **12.2.3 Challenges in Assessing Estrogenic Exposures:**

Studying the link between dietary and environmental estrogens and breast cancer risk faces several challenges:

**Difficulties in Quantifying Exposure:** Accurately measuring an individual's overall exposure to various estrogens is complex. Dietary intake can vary significantly, and environmental exposures can depend on factors like location, occupation, and lifestyle choices.

**Distinguishing Cause from Correlation:** Observational studies often show associations between exposures and outcomes, but they cannot definitively prove causation. Other risk factors for breast cancer might influence the results, making it difficult to isolate the specific impact of estrogens.

**Diversity of Estrogenic Effects:** Different estrogens have varying potencies and may even exhibit anti-estrogenic properties in specific contexts. Understanding the specific effects of different exposures adds another layer of complexity to the research (Vandenberg et al., 2020, Rocha et al., 2021).

### **12.3 Mechanisms of Action: How Estrogenic Exposures Might Influence Breast Cancer Risk:**

The potential link between dietary and environmental estrogens and breast cancer risk hinges on their ability to interact with the body's hormonal system. Here, we delve into the proposed mechanisms by which these exogenous estrogens might influence breast cancer development:

#### **12.3.1 Estrogen Receptor Binding:**

The primary mechanism involves binding to estrogen receptors (ERs). These are proteins located within the cells, particularly in breast tissue.

When natural estrogen binds to ERs, it triggers a cascade of cellular signaling pathways that regulate growth, proliferation, and differentiation. Dietary and environmental estrogens, with their structural similarities to natural estrogen, can also bind to ERs, albeit with varying affinities (Chakraborty et al., 2023).

**There are two main types of ERs:** ER alpha (ER $\alpha$ ) and ER beta (ER $\beta$ ). ER $\alpha$  is considered the primary driver of breast cancer growth, while ER $\beta$  can have opposing effects, potentially acting as a tumor suppressor. The specific type of ER an estrogen binds to and the resulting cellular response can significantly influence its impact on breast cancer risk (Chakraborty et al., 2023).

**Agonistic vs. Antagonistic Effects:** Depending on the specific estrogen and its affinity for ERs, the effects can be either agonistic (mimicking natural estrogen) or antagonistic (blocking the effects of natural estrogen) (Li et al., 2023).

**Xenoestrogens with high ER $\alpha$  affinity:** These xenoestrogens can act as agonists, stimulating similar growth-promoting pathways as natural estrogen, potentially increasing breast cancer risk.

**Phytoestrogens with weaker ER affinity:** These may have mixed effects. At low concentrations, they might act as weak agonists. However, at higher intakes, they might compete with natural estrogen for binding sites, acting as antagonists and potentially offering some protection.

**ER-Independent Pathways:** Some studies suggest that certain xenoestrogens might exert estrogenic effects even through pathways independent of ER binding. These pathways are less understood but could involve interactions with other cellular signaling mechanisms that influence breast cancer development (Lecomte et al., 2017, Wang et al., 2021).

### **12.3.2 Altered Estrogen Metabolism:**

The body naturally breaks down and removes estrogen through a process called metabolism. Certain dietary and environmental estrogens can interfere with this process, potentially leading to elevated circulating levels of estrogen and increased exposure to breast tissue. For instance, some phytoestrogens have been shown to inhibit enzymes involved in estrogen metabolism (Al-Shami et al, 2023).

### **12.3.3 Genotoxic Effects:**

Some xenoestrogens might possess genotoxic properties, meaning they can damage DNA. This damage can lead to mutations that contribute to cancer initiation and progression. The specific mechanisms by which xenoestrogens might induce DNA damage are still being investigated. However, some potential pathways include oxidative stress and direct interaction with DNA repair mechanisms (Calaf et al., 2023).

### **12.3.4 Epigenetic Effects:**

Epigenetics refers to changes in gene expression that don't involve alterations in the DNA sequence itself. These modifications can influence how genes are expressed in breast cells, potentially impacting their growth and differentiation.

Some studies suggest that exposure to certain xenoestrogens might lead to epigenetic modifications that could influence breast cancer risk. These modifications can involve methylation patterns on DNA or alterations in histone modifications, both of which can regulate gene expression (Wang et al., 2021, Calaf et al., 2023).

### **12.3.5 Timing and Duration of Exposure:**

The timing and duration of exposure to estrogenic compounds are also critical factors. Breast tissue is particularly sensitive to hormonal influences during critical developmental stages, such as puberty and pregnancy. Exposure to estrogens during these periods might have a more significant impact on breast cancer risk compared to exposure later in life.

Additionally, the cumulative effects of long-term exposure to low levels of various estrogens might also play a role in breast cancer development, although this requires further investigation (Calaf et al., 2023, Wang et al., 2021).

### **12.3.6 Individual Susceptibility:**

Individual susceptibility to the effects of estrogens can vary depending on genetic factors and overall hormonal milieu. Certain genetic polymorphisms might influence how efficiently the body metabolizes estrogens or how cells respond to estrogen signaling, potentially making some individuals more susceptible to the effects of exogenous estrogens. Additionally, pre-existing risk factors for breast cancer, such as family history or mutations in BRCA genes, might interact with exposure to estrogens, further influencing risk (Sun et al., 2015).

## **12.4 Current Understanding and Controversies: Navigating the Uncertainties:**

The research on the link between dietary and environmental estrogens and breast cancer risk paints a complex picture, with both intriguing possibilities and lingering uncertainties. Here's a deeper dive into the key findings and ongoing debates:

### **12.4.1 Key Findings: A Landscape of Possibilities and Inconsistencies:**

**Dietary Estrogens:** The story of phytoestrogens and breast cancer risk remains a work in progress. While some studies suggest no association, others hint at a possible protective effect at moderate intake levels, particularly for soy isoflavones. However, the research is not entirely conclusive (Mauny et al., 2022, Rocha et al., 2021). Factors like the type of soy product consumed, duration of intake, and an individual's overall hormonal milieu might influence the effects. Additionally, some studies raise concerns about potential negative effects of high-dose isoflavone supplements, highlighting the importance of a balanced dietary approach (Rocha et al., 2021).

**Mycoestrogens:** These estrogenic compounds produced by molds pose potential concerns. Some studies suggest a link between exposure to mycoestrogens, like zearalenone, and an increased risk of certain hormone-related cancers, including breast cancer. However, the specific mechanisms of action and the strength of this association require further investigation. Additionally, factors like the extent of dietary exposure and individual susceptibility play a role, making it difficult to establish definitive cause-and-effect relationships (Fontaine et al., 2020, Paterni et al., 2017).

**Environmental Estrogens:** Studies investigating xenoestrogens show mixed results, adding to the complexity of the picture. Some studies suggest a possible association between exposure to certain xenoestrogens, like BPA, and increased breast cancer risk. BPA is a common chemical found in some plastics and has been shown to exhibit weak estrogenic activity. However, other studies haven't found a clear link, raising questions about the generalizability of findings and the potential influence of other factors. Additionally, research on other xenoestrogens, such as phthalates found in some personal care products, is ongoing, with the potential impact on breast cancer risk still being explored (Wang et al., 2021, Stillwater et al., 2020).

### **12.4.2 Moving Forward: A Call for Collaborative Research:**

Despite the uncertainties, exploring potential preventative measures to reduce exposure to environmental estrogens is crucial. Additionally, focusing on research efforts that address the current limitations can shed light on this complex relationship. Here are some key areas for further exploration:

**Understanding Specific Estrogens:** Differentiating the effects of various dietary and environmental estrogens requires a deeper understanding of their specific interactions with the body's hormonal system. Research investigating the mechanisms of action for different estrogens, including their binding affinities to ERs, potential for epigenetic modifications, and interactions with other signaling pathways, is essential (Yuan et al., 2023).



**Long-Term Studies:** Longitudinal studies that track individuals over time and account for various factors influencing breast cancer risk can provide more robust evidence. These studies can help elucidate the potential long-term effects of chronic low-level exposure to various estrogens and their cumulative impact on breast cancer development (Kuan et al., 2023).

**Genetic Susceptibility:** Investigating the role of genetics in individual susceptibility to the effects of estrogens is also crucial. Certain genetic polymorphisms might influence how efficiently the body metabolizes estrogens or how cells (Sun et al., 2015).

### **12.5 Potential Preventative Measures: Empowering Individuals with Choices:**

Despite ongoing debates, exploring potential preventative measures to reduce exposure to environmental estrogens is crucial.

Here are some possibilities, along with considerations for a more comprehensive approach:

**Reducing Reliance on Plastics:** Choosing alternatives to BPA-containing plastics, especially for food storage and hot beverages, can be a starting point. However, the landscape of plastic alternatives is constantly evolving. Research is ongoing to identify truly safe and sustainable alternatives that are free from potential endocrine disruptors. Additionally, focusing on proper recycling practices and advocating for regulations on safer plastics in manufacturing can be part of a broader solution (Akash et al., 2023).

**Dietary Choices:** While research on phytoestrogens remains inconclusive, maintaining a balanced diet rich in fruits, vegetables, and whole grains is recommended for overall health. This dietary pattern is generally associated with numerous health benefits, including promoting a healthy gut microbiome, which can influence estrogen metabolism. Consulting with a registered dietitian can help create a personalized dietary approach that incorporates these considerations while also addressing individual taste preferences and cultural dietary practices (Chen et al., 2023).

**Personal Care Products:** Considering fragrance-free or paraben-free options for personal care products might be a way to minimize exposure to potential xenoestrogens found in some cosmetics and lotions. However, navigating the often-complex ingredient lists on personal care products can be challenging. Supporting initiatives that promote transparency in product labeling and advocate for the use of safer ingredients in the cosmetics industry can be a valuable approach (Vandenberg et al., 2020).

**Healthy Lifestyle Habits:** Maintaining a healthy weight, engaging in regular physical activity, and limiting alcohol consumption are well-established strategies for reducing breast cancer risk. These lifestyle choices can also positively influence overall hormonal balance and potentially mitigate the effects of estrogen exposures. Encouraging access to affordable and inclusive physical activity options, promoting healthy eating habits through community-based programs, and addressing social determinants of health that influence lifestyle choices are crucial aspects of promoting healthy lifestyles for breast cancer prevention (Satpathi et al., 2023).

## **12.6 Conclusion:**

The relationship between estrogens and breast cancer risk presents a compelling yet intricate puzzle for scientific inquiry. While dietary and environmental estrogens hold potential for influencing breast cancer development, the current research landscape reveals a nuanced picture with lingering uncertainties. On the one hand, the inconclusive findings regarding phytoestrogens highlight the need for further exploration of their potential benefits and risks. On the other hand, the possibility of xenoestrogens contributing to breast cancer development underscores the importance of minimizing exposure to these environmental contaminants.

Public health initiatives play a crucial role in navigating this complexity. Empowering individuals with actionable steps, such as reducing reliance on BPA-containing plastics, maintaining healthy dietary patterns, and prioritizing healthy lifestyle habits, can contribute to overall well-being and potentially mitigate the effects of estrogen exposures. However, these efforts must be coupled with continued research endeavors.

Long-term studies investigating the long-term effects of chronic low-level exposure and the role of genetic susceptibility hold immense promise for unraveling the mechanisms at play. Additionally, fostering open and honest risk communication through various channels is essential for empowering individuals and informing policy decisions.

Ultimately, a multifaceted approach that integrates scientific research, public health initiatives, and collaborative efforts involving researchers, healthcare professionals, policymakers, and the public is key to addressing the complexities of the estrogen-breast cancer link. By working together, we can navigate the current uncertainties and strive towards a future with improved breast cancer prevention strategies. This future holds the potential for not only reducing unnecessary exposures but also developing targeted interventions that address individual risk profiles and pave the way for a more personalized approach to breast cancer prevention.

## **12.7 References:**

1. Akash, M. S. H., Rasheed, S., Rehman, K., Imran, M., & Assiri, M. A. (2023). Toxicological evaluation of bisphenol analogues: preventive measures and therapeutic interventions. *RSC advances*, 13(31), 21613-21628.
2. Al-Shami, K., Awadi, S., Alsheikh, A. M., Al-Sharif, S., Ala'Bereshy, R., Al-Eitan, S. F., ... & Al Zoubi, M. S. (2023). Estrogens and the risk of breast cancer: A narrative review of literature. *Heliyon*.
3. Awuchi, C. G., Ondari, E. N., Nwozo, S., Odongo, G. A., Eseoghene, I. J., Twinomuhwezi, H., ... & Okpala, C. O. R. (2022). Mycotoxins' toxicological mechanisms involving humans, livestock and their associated health concerns: A review. *Toxins*, 14(3), 167.
4. Bilal, M., Rizwan, K., Adeel, M., Barceló, D., Awad, Y. A., & Iqbal, H. M. (2022). Robust strategies to eliminate endocrine disruptive estrogens in water resources. *Environmental Pollution*, 306, 119373.

5. Calaf, G. M., Ponce-Cusi, R., Aguayo, F., Muñoz, J. P., & Bleak, T. C. (2020). Endocrine disruptors from the environment affecting breast cancer. *Oncology letters*, 20(1), 19-32.
6. Chakraborty, B., Byemerwa, J., Krebs, T., Lim, F., Chang, C. Y., & McDonnell, D. P. (2023). Estrogen receptor signaling in the immune system. *Endocrine reviews*, 44(1), 117-141.
7. Chen, Z., Qian, F., Hu, Y., Voortman, T., Li, Y., Rimm, E. B., & Sun, Q. (2023). Dietary phytoestrogens and total and cause-specific mortality: Results from 2 prospective cohort studies. *The American Journal of Clinical Nutrition*, 117(1), 130-140.
8. Desmawati, D., & Sulastri, D. (2019). Phytoestrogens and Their Health Effect. *Open Access Macedonian Journal of Medical Sciences*, 7(3), 495–499.
9. Domínguez-López, I., Yago-Aragón, M., Salas-Huetos, A., Tresserra-Rimbau, A., & Hurtado-Barroso, S. (2020). Effects of dietary phytoestrogens on hormones throughout a human lifespan: A review. *Nutrients*, 12(8), 2456.
10. Fontaine, C., Morfoisse, F., Tatin, F., Zamora, A., Zahreddine, R., Henrion, D., ... & Garmy-Susini, B. (2020). The impact of estrogen receptor in arterial and lymphatic vascular diseases. *International Journal of Molecular Sciences*, 21(9), 3244.
11. Giaquinto, A. N., Sung, H., Miller, K. D., Kramer, J. L., Newman, L. A., Minihan, A., ... & Siegel, R. L. (2022). Breast cancer statistics, 2022. *CA: a cancer journal for clinicians*, 72(6), 524-541.
12. Gompel, A., Fain, O., Boccon-Gibod, I., Gobert, D., & Bouillet, L. (2020). Exogenous hormones and hereditary angioedema. *International Immunopharmacology*, 78, 106080.
13. Gupta, R., Kumar, P., Fahmi, N., Garg, B., Dutta, S., Sachar, S., ... & Vimalaswaran, K. S. (2020). Endocrine disruption and obesity: A current review on environmental obesogens. *Current Research in Green and Sustainable Chemistry*, 3, 100009.
14. Ionescu, V. S., Popa, A., Alexandru, A., Manole, E., Neagu, M., & Pop, S. (2021). Dietary phytoestrogens and their metabolites as epigenetic modulators with impact on human health. *Antioxidants*, 10(12), 1893.
15. Kinkade, C. W., Rivera-Núñez, Z., Gorczyca, L., Aleksunes, L. M., & Barrett, E. S. (2021). Impact of fusarium-derived mycoestrogens on female reproduction: A systematic review. *Toxins*, 13(6), 373.
16. Kuan, C. C., Kuo, W. H., Chang, S. H., & Sun, H. F. (2023). A longitudinal study on the changes in the self-efficacy of breast cancer patients during adjuvant chemotherapy. *Nursing Open*, 10(5), 2912-2919.
17. Lecomte, S., Demay, F., Ferrière, F., & Pakdel, F. (2017). Phytochemicals Targeting Estrogen Receptors: Beneficial Rather Than Adverse Effects? *International Journal of Molecular Sciences*, 18(7), 1381.
18. Li, J., Yu, J., Zou, H., Zhang, J., & Ren, L. (2023). Estrogen receptors-mediated health benefits of phytochemicals: A review. *Food & Function*.
19. Maharjan, C. K., Mo, J., Wang, L., Kim, M. C., Wang, S., Borchering, N., ... & Zhang, W. (2021). Natural and synthetic estrogens in chronic inflammation and breast cancer. *Cancers*, 14(1), 206.
20. Mauny, A., Faure, S., & Derbré, S. (2022). Phytoestrogens and Breast Cancer: Should French Recommendations Evolve? *Cancers*, 14(24), 6163.
21. Miziak, P., Baran, M., Błaszczak, E., Przybyszewska-Podstawka, A., Kałafut, J., Smok-Kalwat, J., ... & Stepulak, A. (2023). Estrogen Receptor Signaling in Breast Cancer. *Cancers*, 15(19), 4689.

22. Paterni, I., Granchi, C., & Minutolo, F. (2017). Risks and benefits related to alimentary exposure to xenoestrogens. *Critical reviews in food science and nutrition*, 57(16), 3384-3404.
23. Paterni, I., Granchi, C., & Minutolo, F. (2017). Risks and benefits related to alimentary exposure to xenoestrogens. *Critical Reviews in Food Science and Nutrition*, 57(16), 3384–3404.
24. Rietjens, I. M. C. M., Louisse, J., & Beekmann, K. (2017). The potential health effects of dietary phytoestrogens. *British Journal of Pharmacology*, 174(11), 1263–1280.
25. Rocha, P. R. S., Oliveira, V. D., Vasques, C. I., Dos Reis, P. E. D., & Amato, A. A. (2021). Exposure to endocrine disruptors and risk of breast cancer: a systematic review. *Critical reviews in oncology/hematology*, 161, 103330.
26. Satpathi, S., Gaurkar, S. S., Potdukhe, A., Wanjari, M. B., & Wanjari, M. (2023). Unveiling the role of hormonal imbalance in breast cancer development: A comprehensive review. *Cureus*, 15(7).
27. Stillwater, B. J., Bull, A. C., Romagnolo, D. F., Neumayer, L. A., Donovan, M. G., & Selmin, O. I. (2020). Bisphenols and risk of breast cancer: a narrative review of the impact of diet and bioactive food components. *Frontiers in nutrition*, 7, 581388.
28. Sun, M. Y., Du, H. Y., Zhu, A. N., Liang, H. Y., de Garibay, G. R., Li, F. X., Li, M., & Yang, X. X. (2015). Genetic polymorphisms in estrogen-related genes and the risk of breast cancer among Han Chinese women. *International Journal of Molecular Sciences*, 16(2), 4121–4135.
29. Thapa, A., Horgan, K. A., White, B., & Walls, D. (2021). Deoxynivalenol and zearalenone—synergistic or antagonistic agri-food chain co-contaminants? *Toxins*, 13(8), 561.
30. Vandenberg, L. N., Najmi, A., & Mogus, J. P. (2020). Agrochemicals with estrogenic endocrine disrupting properties: Lessons Learned? *Molecular and cellular endocrinology*, 518, 110860.
31. Wang, J., Huang, Y., Wang, S., Yang, Y., He, J., Li, C., ... & Martyniuk, C. J. (2021). Identification of active and inactive agonists/antagonists of estrogen receptor based on Tox21 10K compound library: Binomial analysis and structure alert. *Ecotoxicology and Environmental Safety*, 214, 112114.
32. Wang, X., Ha, D., Yoshitake, R., Chan, Y. S., Sadava, D., & Chen, S. (2021). Exploring the biological activity and mechanism of xenoestrogens and phytoestrogens in cancers: emerging methods and concepts. *International journal of molecular sciences*, 22(16), 8798.
33. Yuan, M., Chen, S., Zeng, C., Fan, Y., Ge, W., & Chen, W. (2023). Estrogenic and non-estrogenic effects of bisphenol A and its action mechanism in the zebrafish model: An overview of the past two decades of work. *Environment international*, 107976.
34. Zimmermann, L., Bartosova, Z., Braun, K., Oehlmann, J., Völker, C., & Wagner, M. (2021). Plastic products leach chemicals that induce in vitro toxicity under realistic use conditions. *Environmental science & technology*, 55(17), 11814-11823.