

1. Effect of Concurrent Strength and Endurance Training on Selected Endurance Parameters

P. Senthil, S. Arul

Department of Physical Education and Sports Sciences,
Annamalai University, India.

Abstract:

The purpose of this study was to find out the effect of concurrent strength and endurance training on selected endurance parameters namely Cardio-respiratory endurance and Resting pulse rate. To achieve this purpose, thirty students from the Department of Physical Education and Sports Sciences, Annamalai University, Annamalai Nagar, Tamil Nadu and India were selected as subjects at random. Their age ranged between eighteen to twenty-two years. The selected subjects were divided into two equal groups of fifteen each namely concurrent strength and endurance training group and control group.

The experimental group has undergone twelve weeks of concurrent strength and endurance training, whereas the control group maintained their daily routine activities and no special training was given. The subjects of the two groups were tested on selected variables namely Cardio-respiratory endurance and Resting pulse rate standardized tests, namely Cooper's 12-minute walk/run test and counting on radial artery/minute at prior and immediately after the training period. The collected data were statistically analyzed through analysis of covariance (ANCOVA) to find out the significant difference, if any between the groups. The .05 level of confidence was fixed to test the level of significance which was considered as an appropriate. The results of the study showed that there were significant differences exist between concurrent strength and endurance training group and control group. Also, concurrent strength and endurance training group showed significant improvement on Cardio-respiratory endurance and Resting pulse rate compared to the control group.

Keywords:

Concurrent Strength and Endurance Training % Cardio-Respiratory Endurance and Resting Pulse Rate % Analysis of Covariance (ANCOVA)

1.1 Introduction:

The sequential exercise regime is referred to as ‘concurrent training’. Concurrent strength and endurance training is undertaken by numerous athletes in various sports in an effort to achieve adaptations specific to both forms of training. Literature findings to date, investigating the neuromuscular adaptations and performance improvements associated with concurrent strength and endurance training (referred to as concurrent training) have produced inconsistent results. Some studies have shown that concurrent training inhibit the development of strength and power, but does not affect the development of aerobic fitness when compared to either mode of training alone. Other studies have shown that concurrent training has no inhibitory effect on the development of strength and endurance. Cardio-respiratory endurance of the ability to carry a work load for a relatively prolonged period [1].

The endurance is that enables the heart, blood vessels and lungs to receive oxygen and take it to the muscular and to do it as often and effortless as possible [2].

Strength and endurance training regimes represent and induce distinctly different adaptive responses when individually performed. Typically, strength-training programs involve large muscle group activation of high- resistance low-repetition exercises to increase the force output ability of skeletal muscle [3].

In contrast, endurance-training programs utilize low-resistance, high-repetition exercises such as running or cycling to increase maximum O₂ uptake (VO₂ max). Accordingly, the adaptive responses in skeletal muscle to strength and endurance training are different and sometimes opposite [4].

Strength training has been reported to cause muscle fiber hypertrophy, associated with an increase in contractile protein, which contributes to an increase in maximal contractile force [3].

Strength training also reduces mitochondrial density and suppresses oxidative enzymes activity which can cause impede endurance capacity, but has minimal impact on capillary density or in the conversion of muscle fibre types from fast twitch (type II fibres) to slow twitch (type I fibres).

In contrast, endurance training usually causes little or no muscle fibre hypertrophy, but it does induce increases in mitochondria content, citric acid enzymes, oxidative capacity and the possibility of muscle fibre conversion from fast twitch to slow twitch. Many competitive endurance athletes incorporate resistance training into their training in a hope to improve endurance performance. However, as previously mentioned adaptations to exercise are generally considered to be specific to the training type of stimulus [5]. Although, many adaptations are specific to the type of training, some changes that occur with resistance training could influence endurance performance, which include: muscle fibre transformations and muscle fibre (type I) hypertrophy, which may alter fibre recruitment patterns and help prevent muscle fatigue, as less motor units need to be activated for the same work load [6].

The purpose of this study was to investigate the effect of concurrent strength and endurance training on selected endurance parameters among physical education students of Annamalai University, Chidambaram, Tamil Nadu and India, on age group ranged between eighteen to twenty-two

1.2 Materials and Methods:

The subjects this study was randomly selected from Annamalai University, Chidambaram. A total of thirty subjects was selected at random and they were divided into two equal groups that are experimental and control groups, each group

consist of fifteen subjects, their age ranged from eighteen to twenty-two. The experimental group has undergone concurrent strength and endurance training such as incline leg press, bench press, jump to box, jump from box, alternate leg squat, alternate leg hops in place, jogging and cycling three days in a week for the period of twelve weeks and the control group was not undergoing concurrent strength and endurance training. To find out the Cardio-respiratory endurance and Resting pulse rate the investigator conducted the Cooper's 12-minute walk/run test and counting on radial artery/minute.

Testers' competency, subject reliability and reliability of tests were established by using test and retest method and the reliability coefficient were found to be satisfactory high.

The data were analyzed using analysis of covariance (ANCOVA) for determine the effect of concurrent strength and endurance training on selected physiological variables of Cardio-respiratory endurance and Resting pulse rate of students of the department of physical education and sports sciences, Annamalai University, Annamalai Nagar, Tamil Nadu and India on the age group between eighteen to twenty-two.

1.2.1 Selection of Variables and Tests:

The research scholar reviewed the available scientific literature pertaining to available the present study, the following variables were selected

Table 1.1: Selection of Variables and Tests

Sr. No.	Variables	Test Items
1.	Cardio-respiratory endurance	Cooper's 12-minute walk/run test
2.	Resting pulse rate	Counting on radial artery/minute

Statistical Analysis: Data were collected from the both groups, the selected variables on Cardio-respiratory endurance and Resting pulse rate was statistically examined by using the “F” ratio used to find out the significance difference between experimental group and control group, the level of significance was fixed at 0.05 level of confidence.

1.3 Results:

Within the limitations of the study the following conclusions appeared and justified as per the result obtained. The mean difference of the criterion measures for the control and experimental groups presented in tables.

Table 1.2 shows that the adjusted post-test means of concurrent strength and endurance training group and control group are 52.63 and 50.79 respectively. The obtained ‘F’ ratio of 87.676 for adjusted post-test means is more than the table value of 4.21 for df 1 and 27 of .05 level. The result of the study indicated that there was a significant difference between the concurrent strength and endurance training group and control group on cardio-respiratory endurance. The mean difference of the criterion measures for the control and experimental groups presented in tables.

Table 1.3 shows that the adjusted post-test means of concurrent strength and endurance training group and control group are 72.67 and 75.00 respectively. The obtained ‘F’ ratio of 9.394 for adjusted post-test

Table 1.2: One way analysis of covariance (ANCOVA) for pretest and post test scores of Cardio-respiratory endurance of control and experimental group.

Test	CSE Group	Con Group		Sum of Squares	Df	Ms	F
Pre-Mean	50.09	51.48	B	10.24	1	10.24	1.387
SD	2.255	3.025	W	206.776	28	7.385	

Test	CSE Group	Con Group		Sum of Squares	Df	Ms	F
Post Mean	55.174	50.311	B	193.918	1	193.918	58.426
SD	1.246	2.371	W	92.934	28	3.319	
Adjusted Post Mean	52.63	50.79	B	216.31	1	216.31	87.676
			W	66.612	27	2.467	

Table value of (1, 27) at 0.05 level was 4.21.

Table 1.3: One way analysis of covariance (ANCOVA) for pretest and post test scores of Resting pulse rate of control and experimental group.

Test	CSE Group	Con Group		Sum of Squares	Df	Ms	F
Pre-Mean	74.4	74.67	B	3.33	1	3.33	0.3846
SD	3.397	2.743	W	242.67	28	8.67	
Post Mean	70.93	75.33	B	90.133	1	90.133	10.073
SD	2.520	3.132	W	250.533	28	8.95	
Adjusted Post Mean	72.67	75.00	B	86.86	1	86.86	9.394
			W	249.647	27	9.25	

Table value of (1, 27) at 0.05 level was 4.21.

means is more than the table value of 4.21 for df 1 and 27 of .05 level. The result of the study indicated that there was a significant difference between the concurrent strength and endurance training group and control group on resting pulse rate.

1.4 Discussion:

By understanding a systemic training at a level above normal, a variety of strength and endurance adaptations take place in the body that makes it function a more effectively. Strength training develops strength parameters and endurance training develops endurance performances separately. Maximal strength training improves aerobic endurance performance [7] but endurance training did not improve the strength performances. Combined strength and endurance training on the compatibility on improving both VO_2 peak and strength performance simultaneously [8]. Concurrent strength and endurance training enhances strength and endurance measures [9]. There was no effect on the sequence of training either endurance training or strength training given as initial one [10].

The study done by the investigator is a unique one and is in consonance with the studies already done by the following authors given in references. The result of the study indicated that systematic and regular level of cardio respiratory endurance and resting pulse rate also to promote performance in any kind of sports activities.

This change or modification on different variables of motor fitness is largely depends on the type of activity, this study clearly indicates that there is a significant difference on cardio respiratory endurance and resting pulse rate under concurrent strength and endurance training program when compared with control group.

This may be the fact that the lungs form the chief oxygenating apparatus in the body it will be realized how very necessary if keep them in a healthy condition for carrying out their work efficiently. degrades up on the phenomena are very closely related to Cardio-respiratory fitness as well as Resting pulse rate, degrades up on the phenomena are very closely related to each other. Base on the demand of oxygen and nutrients values in the muscles, duo to physical and mental load, the supply of oxygen as well as nutrients is gradually improved through the rate of blood circulations in the working muscles.

Concurrent strength and endurance training such as incline leg press, bench press, jump to box, jump from box alternate leg squat, alternate leg hops in place, jogging and cycling exercises might have been successfully improved when compare with the control group. This may be due to the practices of (12 weeks) concurrent strength and endurance training program. This study also proved that the cardio respiratory endurance as well as the resting pulse rate significantly improved when compared with control group. This may due to the fact that concurrent strength and endurance exercises program influenced as them.

1.5 Conclusion:

The finding of this study indicated that the effects of concurrent strength and endurance training showed significantly improved the Cardio-respiratory endurance and reduced Resting pulse rate when compared with control group.

1.6 References:

1. Lawrence, E More House and Angutus, T. Miller, 1968. *Physiological of Exercise*, (St. Louis: The C.V. Mosby Company, pp: 67.
2. France Walk Field, Dorothy HawKing and John M. Cooper, 1973. *Track and Field Fundamentals for Girls and Women*, (Saint Louis: The C.V. Mosby Company, pp: 22.
3. Sale, D.G., J.D. MacDougal, I. Jacobs and S. Garner, 1990. Interaction between concurrent strength and endurance training. *Journal of Applied Physiology*, 68: 260-270.
4. Tanaka, H and T. Swensen, 1998. Impact of resistance training on endurance performance: A new form of cross-training? *Sports Medicine*, 25: 191-200.
5. Nelson, A.G., D.A. Arnall, S.F. Loy, L.J. Silvester and R.K. Conlee, 1990. Consequences of combining strength and endurance regimens. *Physical Therapy*, 70: 287-294.

6. Bishop, D. and D. Gjenkins, 1999. The effects of strength training on endurance performance and muscle characteristics. *Medicine and Science in Sports and Exercise*, 31: 886-891.
7. Hoff, J., A. Gran and J. Helgerud, 2002. Maximal strength training improves aerobic endurance performance, *Scandinavian Journal of Medicine & Science in Sports*, 12(5): 288-095.
8. McCarthy, J.P., J.C. Agre, B.K. Graf, M.A. Pozniak and A.C. Vailas, 1995. Compatibility of adaptive responses with combining strength and endurance training. *Medicine & Science in Sports & Exercise*, 27(3): 429-36.
9. Davis, W.J., D.T. Wood, R.G. Andrews, L.M. Elkind and W.B. Davis, 2008. Concurrent training enhances athletes' cardiovascular and cardiorespiratory measures. *The Journal of Strength & Conditioning Research*, 22(5): 1503-14.
10. Collins, M.A. and T.K. Snow, 1993. Are adaptations to combined endurance and strength training affected by the sequence of training? *Journal of Sports Sciences*, 11(6): 485-91.