

8. In Vitro Shoot Induction from Stem Node Explants of *Trichosanthes Cucumerina* (L)

Mandalaju Venkateshwarlu

Department of Botany,
Kakatiya University,
Warangal, Telangana State, India.

Abstract:

The effect of BAP and NAA + Kn combination on In vitro shoot regeneration and elongation was analysed in shoottip explants. The percentage of response was increased upto 2.0mg/L to 3.0 mg/L Kn and later gradually decreased at high concentrations. Plantlet regeneration through Cotyledon and shoottip explants and streptomycin resistant plantlet using in vitro mutagenesis was also developed in S. Surattence (Muukherjee et al 2011, Swamy et al 2005, Nasar et al 1997) plant regeneration callus induction.

*In vitro regeneration in the *Trichosanthes cucumerina* (L) study of multiple shoots MS Containing media with different and combinations of growth regulators. The effect of plant growth regulators on rooting was investigated 4-6 weeks after culture on MS medium (Subashri et al 2014: Devendra et al 2011: Verma 1996: Sunderraj et al 1989). Due to over exploitation for high medicinal values and destruction of the habit this Sunderraj et al 1989, Nasir et al 1997) meristem shoottip cultures hence there is need for ex situ conservation through tissue culture methods. The stem node explants were inoculated on MS medium supplemented with various cytokinins i.e., BAP and NAA. Addition of BAP at 3.0 mg/L concentration or NAA at 3.0 mg/L to the MS basal medium, induced regeneration from the stem node explants. Micro propagation involves multiplication of genetically identical individual by asexual reproduction within a short span of time with tremendous potential for the production of high-quality plant-based medicines (Murch et al., 2000). Multiple shoot induction was achieved in one of the important medicinal plants of *Trichosanthes Cucumerina* (L) MS medium supplemented with 1.0 mg/l BAP + 2.0 mg/l NAA and 3.0 mg/l L-Glutamicacid was found to be optimum to induce shoots. The present study established reliable and reproducible protocol for rapid multiple shoot induction from stem node explants of *Trichosanthes cucumerina* using different concentrations and combinations of Cytokinins.*

Keywords:

Stem node, NAA BAP, Trichosanthes Cucumerina, Shoot induction

8.1 Introduction:

Stem node explants of *Trichosanthes Cucumerina* (L) on MS medium fortified with plant growth regulators along with coconut milk and Amino acids. The plants of Cucurbitaceae suffer from several diseases including the water melon mosaic virus (Greber, 1978), Cucumber green mottle mosaic virus (Nijssen, 1984) and *Trichosanthes Cucumerina* (L) also suffers from downy and powdery mildews which seriously limits the crop production.

Auxiliary buds from pumpkin were reported by Jelaska (1972). *In Vitro* Regeneration of plants via Somatic embryo genesis has much potential for plant propagation and gene transfer in Soybean somatic embryo have been obtained from cultured immature cotyledons Soybean from embryogenic callus line derived from the shoot tip explants (Parvathi, Venkateshwarlu M-2018) *In Vitro* Regeneration Via somatic embryo genesis has drawn more attention than other methods because it can produce a large number of plants in a relatively short time (Walker *et. al* 2001) The number of protoplasts showed increase during shorter treatment time and reaches a peak at 4-5 hours of after this many reports of soybean somatic embryogenesis were published (Pathak. *et.al* 2014)

The development of protoplast systems has increased the plants for use in both Biochemical and genetic research. Ugandhar *et. al* (2011) has been also reported that high amount of Cytokinin and lower number of Auxins is the best combination for Somatic embryogenesis which is in accordance of our study.

8.2 Materials and Methods:

The presence of Cytokinin inducing shoot organogenesis well developed shoot lets from our *Trichosanthes cucumerina* experimental data. The capacity of shoot bud differentiation and shoot proliferation from shoot tip explants of Coconut milk (CM) depended on hormonal variation.

The percentage of response was increased gradually from 2.0 to 3.0 mg/L BAP, the shoots raised *in vitro* 2-3 cm long were cultured on MS medium supplemented with various concentrations multiple shoot formation from shoot apics was obtained on

MS medium supplemented with 1.0mg/L NAA (Deca *et al* 1999 and Anju *et al* 2005) shoot tip explants showed a positive response on MS medium containing BAP (1.0 mg/L-2.0mg/L) combination with NAA (1.0mg/L to 2.0 mg/L) That clearly implies the additive effect of IAA (1.0mg/L to 2.0 mg/L) combination of auxin and Cytokinin favored shoot bud differentiation in many plants. BAP was superior to Kn in inducing high frequency shoot regeneration in many numbers of plants. (Srilatha, Venkateshwarlu *m* 2019) planlet regeneration from Cytoledon culture Soybean All media were adjusted to PH 5-8 before addition of 0.8% agar-agar and auto claved at 121°C and 103K Pg for 20Min Cultures 25x 150nm cultures tubes. MS Basal Medium was supplemented with various plant growth regulators and 3.0% Sucrose.

The PH of the media was adjusted to 5.8, solidified with 0.8% Difco-bacto Agar and Autoclaved at 103.4 KPa or 121°C for 15-20min. A single explant was placed in each culture tube and incubated (at 25± 1°C with a 16h photoperiod under fluorescent light (40-50m² s⁻¹).

Explants with in Vitro multiple shoots proliferated on TD2-containing media were transferred to MS Medium containing different concentrations (BAP 2.0-3.0mg/L+) Multiple shoot intonation from shoot tip explants was observed within 20-25days after inoculation. The effect of different five types of growth regulators on direct plantlet regeneration of tamato from shoot tip explants (T Ugandar & M Venkateshwarlu-2018)

The Result from this study has shown that BAP induced the activation of Totipotency at the stem node explants, which resulted in the formation of multiple shoots.

The stem node segments of 2.0 – 3.0 cm long were cultured and surface sterilized with 0.1% HgCl₂ for 5-7 minutes and rinsed with sterile distilled water. They were cultured on MS medium containing 2.5% sucrose and 0.8% Agar-Agar and different concentrations of BAP, NAA and L-Glutamic acid. The pH of the medium was adjusted to 5.8 and later was autoclaved at 120°C for 17 minutes. Cultures were incubated under 16 hrs, illumination (250 lux) at 25±2°C temperature. Raising the level of BAP (0.5 to 2.0 mg/L) resulted in the increase in the number of shoots from hypocotyls and cotyledon explants of Niger (Nikam and Shitole, 1993). Cotyledonary explants (3-4 weeks old) of different sizes (0.5-1.0mm) were cultured with the abaxial surface in contact with induction Ms Medium consisting (1.0-2.0 mg/L BAP) and (1.0 to 3.0 mg/L) NAA for maturation and plant regeneration.

8.3 Results and Discussion:

The hormonal supplement was selected because it was optimum for callus formation stem node explants of *Trichosanthes cucumerina* developed (60%) with increase in concentration of NAA in MS Medium. The regenerated plants will be useful for constant supply of uniform raw materials for commercial secondary metabolic extraction according to their observation. BAP, Kn were superior for multiple shoot formation has been reported as it was observed in the present investigations. Then the plantlets were transferred to polypots containing pre chamber set at 28°C and 60-70% relative humidity. After 2-4 weeks they were transplanted to polybags containing mixture of soil+sand+manure in 1:1:1 ratio and kept under shade house for a period of 6 days to 4 weeks. Multiple shoot buds were initiated on the callus cultured in MS Medium supplemented with different combinations (BAP, NAA & Kn). The regenerated elongated shoots were transferred to Indole Byutyric acid (IBA) (2.0mg/L-4.0mg/L) for root induction. Sub cutting of callus into fresh medium containing the same concentrations of growth regulates resulted in the emergence of callus. The Stem node cuttings were inoculated on MS basal medium fortified with various cytokinins i.e., BAP and NAA. Coconut water also had a role in triggering the formation of multiple shoots. Raising the level of BAP (3 mg/l to 4 mg/l) resulted in an increase in the percentage of shoots developed from Stem node cuttings.

There was no significant increase in the number of shoots on NAA at low and high concentration. Low concentration of L-glutamic acid (0.5 – 1.0 mg/l, along with BAP (2.0 mg/l, has produced significant mean number of multiple shoots that ranged from 2-3 to 5-6 in both the explants. The mean number of shoots developed on the explants ranged from 1-4 to 2-3 by the addition of different concentrations of BAP and NAA. The number of shoots developed on the explants ranged from 1-4 to 2-3 by the addition of BAP at a concentration of 2.0 mg/l or NAA at 2.5 mg/l. (Plate – 8.1, Table –8.1). MS medium fortified with 1.0 mg/l BAP or 2.0 mg/l L-Glutamic acid also induced shoot buds on Stem node explants. Addition of NAA failed to produce many shoots but enlarged the stem node segments. Lower levels of coconut milk (6, 12%) induced callus formation. The results from study have shown the initiation of shoot buds and formation of multiple shoots from different explants i.e. Stem node cuttings of *Trichosanthes Cucumerina* (L). Among all explants used Stem node segments were the best for multiple shoot induction. With an increase in the level of BAP 2.0 – 3.0 mg/l the percentage of explants producing shoots also increased.

Table 8.1: *In Vitro* Shoot induction from Stem node explants of *Trichosanthes Cucumerina* (L).

Growth Regulators	Stem Node explants	
	% frequency of Shoots	Mean No. of Shoots
MS + 0.5 mg/l BAP + 1.0 L-Glutamic acid + IBA	30	Green Callus
MS + 1.0 mg/l BAP + 2.0 L-Glutamic acid + IBA	40	Green Callus
MS + 2.0 mg/l BAP + 3.0 L-Glutamic acid + IBA	35	Callus + shoots (1-3)
MS + 3.0 mg/l BAP + 4.0 L-Glutamic acid + IBA	30	Callus + shoots (4-6)
MS + 1.5 mg/l NAA + 0.5mg/l	25	Green Callus
MS + 1.0 mg/l NAA + 2.0mg/l	22	Green Callus
MS + 2.5 mg/l NAA + 2.5mg/l	20	Callus + shoots (2-4)
MS + 3.0 Mg/l NAA + 3.0mg/l	15	Callus + shoots (1-3)
MS + 3.5 mg/l NAA + 3.0mg/l	10	Callus + shoots (3-6)
CM = Coconut milk water		



Plate 8.1: *In Vitro* Shoot Induction from Stem node explants of *Trichosanthes Cucumerina* (L).

8.4 Conclusion:

Maximum shoot bud differentiation from callus culture was achieved on MS Medium supplemented with BAP, NAA and KN, therefore their MS Medium was designated as multiple shoots. The callus showed maximum number of shoot bus (2-4) The method of repeated transfer of explant is considered to be useful for large scale production of plants, as it avoids isolation and culture of new explants. This is considered as one of the methods to increase the response in explants has suggested that repeated transfer of explants on multiplication media containing cytokinins succeeds in activating the plant materials. The purpose of this work was to study the effect of different concentrations of growth regulators on direct plantlet regeneration of *Trichosanthes cucumerina* from stem node explants. Rooted plantlets were successfully hardened under culture conditions and established in the field conditions.

8.5 References:

1. Anju A and chawla S (2005) Organogenic plant regeneration via callus induction in chickpea (*cicer arictinum* L) Role of genotypes growth regulators and explants Indian Journal of Biotechnology 4: 251-256.
2. Deka AC, Kalita MC and Baruah A (1999) Micropropagation via potent herbal medicinal plant withania Somnifera Environ. Ecology 17(3): 594-596
3. Devendra BN, Srinivas N, Sandeep Reddy a (2011) A high frequency Somatic embryogenesis and plant regeneration in nodal explants culture of *Eclipta alba* L Hassk Ann. Noil res 293): 143-149.
4. Greber, R.S. (1978). Water melon mosoic virus 1 & 2 in: Queensland Cucurbit crops. Asst. J Agri. res 29: 1235-1245.
5. Harris, P. J. C. and Moore, T. H. M. (1989). Nitrogen fixing trees Res Rep 7-12b.
6. Halder, T.E. and Gadgil, V.N (1982). Shoot bud differentiation in long term callus cultures of *Momordica* and *Cucumis* Ind. J. Expt. Boil, **20**: 80-782.
7. Jelaska, S. (1974). Embryogenesis and organogenesis in pumpkin explants. *Physiol. Plant.* **31**: 257-261.
8. Komalavalli, N. and Rao, M.N. (2000). In vitro micropropagation of *Gymneme sylvestre* a rare multipurpose medicinal plant. *Plant Tissue Organ Cult.* **61(2)**: 97-105.
9. Mukherjee P, Varshney A, Johnson TS, Jha TB (2011) *Jatropha curcas* a review on biotechnological status and challenges plant Biotechnol. Rep 5: 197-215.
10. Murch, S.J. Krishna Raj, S. and Saxena, P.K. (2000). Tryptophan is a precursor

- for melatonin and serotonin biosynthesis in in vitro regenerated St. Jhon's wort (*Hypericum perforatum* L. cv. Anthos) Plants, Plant Cell Rep. **19**: 698-704.
11. Morel Georges, (1952). Producing virus free cymbidium Am. Orchid. Soc. Bull. 29: 495-497.
 12. Nasiva Saeed Ysuf Zafer and Kauser A Malik (1997) Simple procedures of Gossypium merstem shoottip cultures plant cell Tissue Org. Cult 51: 201-207.
 13. Nikan, T.D. and Shitole, M.E. (1993). Plant Tissue and Organ Culture **32**: 345.
 14. Nijdsen, A.P.M. den (1984). Attempts to introduce resistance from related species into the cucumber. Proceedings of the third meeting on breeding of cucumber and melons 2-5 July 1984 Poldiv Bulgaria
 15. Parvathi D, Venkateshwarlu M, Dinesh Khannaand Yugandhar T (2018) protoplasts isolation of Soybean Glycine max (L.) Merrill from leaf explants the pharma innovation journal 2018:7(10):661-665.
 16. Pathak N, Tiwans, Mishra MK, Gautam SS (2014) Max (L.) Merr. In proceedings of National Conference on Biotechnology Jabalpur, MP, 2014, 41.
 17. Sounderraj V Telavathi DH and Nijalingappa BHM (1989) Shoottip culture in Dolichos biflorus L Current Sci 58: 1385-1388
 18. Subashri b Pillai YJK (20104) in Vitro regeneration of bacopa monnieri (L) pennet A multipurpose medicinal plant Int. Journal Pharma Sci. 6(4) 559-563
 19. Srilatha T Venkateshwarlu M, Anitha Devi U and Ugender T (2019) Efficient 2,4-D and TD2 assisted Somatic embrogenesis and plant let regeneration in Indian Soybean (Glycine max L) from Cotyledonary explants. The pharma innovation journal 8(4): 322-326.
 20. Swamy N, Ugender t Praveen m R Babu M (2005) induction of Streptomycin resistant plantlets in Solanum Surattense through in Vitro mutagenesis plant Cell. Tissue Org. Cult. 80: 201-207.
 21. T Ugandhar, M Venkateshwarlu (2018) *In Vitro* plantlet regeneration of tamoto (*Solanum Lycoperscan Ruby*) from using five MGLL.CV.Pusa Shoot tip explants Cytokinins J. Indian bot.soc. Vol.97
 22. Verma and kant u (1996) Micropropagation of Embilila officnale (Gaertz) through mature nodal explants. J playtol Res 9(2): 107-109.
 23. Walker Dr, Parrott WA – (2001) Effect of polyethylene glycol and sugar alcohols on soyabean somatic embryo germination and conservation. Plant cell Tissue Organ Cult. 64:55-62
 24. Wehener, T.C., Locky, R.D. (1981). In vitro adventitious shoot and root formation of cultivars andlines of Cucumis sativus L Hort. Sci. **16**: 759-760.

25. Yugandhar T, Venkateshwarlu M, Parvathi D, Shekar GPV, Srilatha T Reddy J.K.- (2011) High frequency Somatic embryogenesis and plantlet regeneration from shoot tip explants of soybean. *Science Research Rep.* 1:146-150